1- This cervix (star) is discovered at 20 wks preg, best ttt is

- a- Cryo-coagulation
- b- Laser coagulation
- c- Diathermy
- d- Cone biopsy
- e None of the above

2- That condition in the cervix, the true statement is

- a- Normally found in pregnancy
- b- Leads to lumbar backache (Sacal)
- e- Should be biopsied __oaluantimete
- d- Best treated by cone biopsy
- e- May lead to infertility

できいでしていた!

The ph is compared of chronic var discharge ?

Lockin-f.

3- Contraindication of that imaging study don't include

- a- Suspected blockage of Fallopian tubes
- b- Suspected acute PIDi Placer Juguist & organis
- c- Suspected intrauterine pregnancy
- d- Premenstrual phase -o Thick endown Polse -Ve (es. 1)
- e- During active bleeding from the aterus ____ extra 1/2:10 /1.

4- DES may lead to all the following except

- a- Patulous internal os
- b- T-shaped uterus
- c- Ectopic pregnancy
- d- Vaginal adenosis 一世 5分 元
- E- Septate merus

5- All the following regarding this test are true except

- a- Should be repeated in high risk population yearly
- b- Suspicious results should be confirmed by colposcopy
- Smear is only taken from the ectocervix (2 Port for 1)
- d- Is to be done for postcoital bleeding
- e- Suspicious specimen is related to HPV 16, 18

6- The result of this test have shown dysplastic cells occupying the deep 1/3 of the epithelium. What is your diagnosis

- Ja- CIN I
- b- CIM 2
- e- CIN 3
- d- CIN 4

7- This stain is done for all of these except

- a- Sexually active female with HPV 6, 11
- b- Sexually active female with HPV 16, 18
- c- Sexually active female esp smokers
- 'd- After subtotal hysterectomy
- _e Routinely in pregnancy

Cides

Sulleion.

8- Which is not associating such organism

- a- Vulval itching
- b- Redness of the vulva
- c- Odorless discharge
- d= Strawberry vaginal walls
- e- Dyspareunia

9- Which is the incorrect statement for that organism

- a- If may be discovered in the vaginal fluid of asymptomatic women
- be It flourish with increased vaginal acidity
- c- Usually associated with malodorous discharge
- d- May be associated with dysuria
- e- Vaginal discharge is typically frothy

10- The incorrect statement regarding that test

- a- Normally the epith of ectoex stains brown
- b- Vag wall stains brown during child bearing period
- c- Areas of columnar metaplasia don't stain brown -- c-
- d= Areas that don't stain brown are definitely malig -
- e- It is complementary test during colposcopy

11- All about this figure is true except

- त्रभी It is seen at mideyele
- b- Indicates estrogenic effect
- c- Positive in PCO at day 21 of cycle
- d- Turns --ve at day 18 of cycle
- e Indicates viscid mucoid cervix

12- This condition leads to all of these except

- a- Recurrent breech presentation
- b- Habitual abortion
- _d Ectopic pregnancy
- e- Menorrhagia -- to

13- All are true about such condition, except a Pelviabdominal swelling is mainly uterine b- Presents as Iry amenorrhea c- The commonest presentation is by retention of urine d- May lead to infertility if ttt is delayed —D Adhesions e- Needs surgical intervention	19-Regarding this swelling a It is the commonest cyst in the vulva b- It is present in the anterior part of labium majus (Post > c- Is mainly caused by obstruction of the duet by infection d- Should be excised if the patient is above 30 years e- The main symptom is throbbing pain
a- May be associated with uterus bicornus bicollus b- May be associated with horse show kidney c- Spasmodic dysmenorrhea could occur — A-Presents by Iry amenorrhea e- Presents by dyspareunia	20- Regarding this swelling, all are true except a- The causative organism is mainly E-coli b- Leads to throbbing pain c- Should be drained d- Is associated with pruritus vulvae (Pain) e- May occur due to gonorrhea
26 wks. Which is not a possible association? a- Recurrent breech b- PTL c- Absent left kidney d- Menorrhagia es Normal pregnancy & normal labor	21- All the following about this lesion is true except a Caused by gonorrhea b- Caused by HPV 6,11 c- May occur in the cervix d- May be treated by chemical cautery e- May be associated with CIN
degree perineal tear. The arrow indicates a structure called a- Bulbo-spongiosus b- Ischio-cavernosus c- Perineal body d- Transverse perineal ms e- None of the abo.e	a- It is an ulcer of the ectocervix b- During pregnancy is best treated by conization c- Cauterization should be the first line therapy d- Pap smear is advisable before therapy e- Commonly it causes pain, dysparuenia and backache 23- Causes of that condition doesn't include
7- All the following statements about such trauma are true except a- May be associated with diabetic mother b- High forceps may lead to it c- Surgical repair is done layer by layer d- Intercourse is avoided for 2-3 years (>>>> +6.)	a- Normal pregnancy b- COC c- ERT d- Acute cervicitis — Silo Borr Psp Carcinoma of cervix

24- Such polyps, all are true except

b- May be due to fibroid

a- Are commonly due to chronic cervicitis

c- Should be removed & sent for histopathology

e- Leads to contact bleeding (Mucous Polyte)

e- May be corrected under regional anesthesia
= epidual

18- Such a condition may be associated with, except

- a- Obstructed labor
- b- Lest occipito-anterior
- JUGI
- d- Precipitate labor (Sages
- e- Ureteric injury

25- This pathology is seen by hysteroscopy, the following should be done, except:

- a- Adhesiolysis via hysteroscopy
- b- Insertion of a balloon
- c- Giving steroids
- --- d= Giving progesterone
 - e- Giving cyclic estrogen & progesterone

26- This discharge is associated with, except:

- a- Acidic ph
- b- COC
- c- Severe soreness in the vulva
- _d-' Good response to metronidazole ___ t> -
- e- Bleeding spots after scrapping

27- This discharge is associated with, except

- a- Typically occurs postmenstrual
- b- May be associated with malodor
- c- Vagina is strawberry
- -d= Cultured on Sabaroud's medium
- e- Its organism may be seen in normal females

28- The arrow shows a condition associated with

- a- Normal vagina
- b- Pruritus & dysparuenia in most cases
- c- Responds well to miconazole
- et- Alkaline vaginal ph
- e- Odorless discharge ? _ m/

29- This condition is associated with all, except

- a- Recurrent lower abdominal pain & discharge
- b- Infertility
- c- Ectopic pregnancy
- d- Chlamydia
- e-1117 6,11 1 (X) ; L.

30- All the following about this slide is true, except

- a. Tubes are better removed in infertile patients
- b- IUCD must be removed before treatment
- c- Suspected more with chlamydial infection
- I Leads to habitual abortion
- e- Leads to ectopic pregnancy



MO(0:0/Piny Chronic

31-1 Pdf to development of prolapse include all except

- a- Congenital weakness of supporting ligaments
- b- Postmenopausal atrophy
- c- Injury during childbirth
- d- Cervical polyps
- e= Ovarian tumors

31-2 Which is the correct statement for that condition

- a- Usually associated with acute local pain _____
- b- More common in nulliparas
- c- Not related to instrumental delivery Tix 'vo Cointing
- d- Not related to postmenopausal changes
- May include sacral backache

3? - Regarding this test all are true except

- a- It may reach up to 15 cm
- b- It turns -ve 3 days after ovulation in Mullerian agenesis
- Tve test means water cervical mucous
- d- +ve test depends on estrogen
- e= +ve test means ovulation

33- All the following is true regarding IVF & ET except : Wild Alect

- 2 Down regulation is done by gonadotrophins Colonsing holon of Controls
- b- Ovarian stimulation may be done by Gn RH
- c- More than one embryo is transferred at a time -- Paradous L

JINLUGE

- d- Success rate is inversely proportional to age
- e-May be done in bilateral cornual block bidica 11.5

34- Which is the incorrect statement regarding that syndrome

- a- It is associated with gonadal failure
- b- There is no secondary sexual ecc
- c- The total number of enromosomes is 45
- d- Estrogen replacement therapy may be used _______.
- E- The ovaries may be polycystic

31:- All these statements about this syndrome is correct except

- a- Usually occurs in obese patients
- b- Is associated with insulin resistance
- c- Follicles are mainly seen subcapsular
- FSII/LII ratio is 3:1
- e- May lead to habitual abortion

49- Engagement of the fetal head is documented when

- a. The fetal head is inside the pelvis
- b- The presenting part is just above the level of ischial spines
- c- The vertex is left OA
- -d- The biparietal diameter pass through the pelvic inlet
- e- The fetal skull is fully flexed

50- All of theses are signs of placental separation except

- a- Uterus comes more globular & harder
- b- Suprapubic bulge
- e- Shortening of the umbilical cord
- d- Gush of blood from the vagina
- e- The fundus raises upwards

51- First trimester abortion may be due to: (100 Co minor est)

- a- Inadequate estrogen production in -t. Porton, 5
- H- Chromosome abnormality of the fetus
- e- Incompetence of the internal cervical os
- d- Maternal diabetes
- e- Cytotoxic drugs

52- Cervical incompetence

- a- Is a common cause of 1st trimester abortion
- b. May be congenital
- e- Is associated with previous instrumental delivery /
- d- is diagnosed at follicular phase HSG
- e- is associated with painful abortion &

53- In ectopic pregnancy:

- a- Bleeding precedes pain (Sal'
 - .5- Shoulders tip pain is an important symptom
- c- The isthmus of the tube is the commonest site of implantation (Arr Pullo)
- Let The incidence is greater in women with IUCD
- e- Ultrasonic scan is of no help in diagnosis

54- Which is the incorrect statement for that case

- 12-HCG is doubled every 48 hours
- Laparoscopy is the traditional standard method to confirm diagnosis
- Les Tubal pregnancy commonly presents with abdominal pain
- , d-Medical conservative III is an option on its by Ulbs < 200
- History of previous salpingitis is relevant

y last technologymonianthis with intent own. No. b.

55- The following statements regarding V.mole are correct except

- a- Usually have female karyotype
- b- May be complicated by thyrotoxicosis
- c- May be complicated by ovarian cyst
- d=α-seto-protein is a good marker for disease post-evacuation size G
- e- Is more prevalent in the advanced age group

Indications for a methotrexate chemotherapy following evacuation of a VM usually include except

- a- a rise in HCG titers
- b- plateau HCG titers for 3 successive weeks
- c- failure of HCG titers to return to normal 8 wks after evacuation
- d- appearance of liver or brain metastases
- rapid disappearance of theca lutein cysts (which

57-35 yrs old, G3P2 presented at 8 wks with mild vag bleeding for 3 days. She passes fleshy tissues per vagina presented on the slide. Further investigations don't include 11. 000 0

- a- Chest X-ray
- 13- Amniocentesis
- é- Serum HCG levels
- d- Pelvic U/S
- 58- Correlate
 - a) P.previa with severe bleeding & immature fetus (4)
 - b) P.previa centralis accidentally discovered at 37 wks (3)
 - (1) Mild placental abruption, no fetal distress at 33 wks
- e) Placental edge 3 cm from ex internal os, presenting in labor
 - . Conservative treatment
 - 2. Albowance or trial of vaginal delivery
 - 3. Elective cesarean section
 - 4. Emergency cesarean section
 - 5. Cesarean hysterectomy

59- As regards P. previa, the following are true

- a) It is manifested by painless bleeding
- b) The initial lige, is usually fatal
- c) It may predispose to PPHge
- d) Its incidence is unaffected by parity
- e) Its incidence decreases with maternal age

60-	The following are risk factors for placental abruption except
	a) Smoking
	b) Folic acid deficiency
	c) Preeclampsia
	History of threatened abortion " " " " " " " .
	e) Previous placental abruption $\frac{1000}{1000}$
<i>~</i> 9 9	This hematoma was found on inspection of the maternal side
01- 1	of the placenta after delivery. The risk factors for this
	condition don't include
	$(1-1)[\cdot]$
	b- Smoking to
	fron deficiency anemia $4 \times 100 \times 5 \times 1000 \times 1000$
	d- Chronic hypertension
	e- External trauma to the abdomen
62-	A diagnosis of severe PET in 37 wks with BPr 160/110 is
,	supported by
	a- Urine output of 1000 ml/24 hrs
	b- That physical sign on the slide - D Pital 13 555 Ands
11	e Epigastric pain Contra So.
	d- BPr of 160/110 at 8 wks in the same gestation 🗡
	e- Λ parity $> 5 \gamma \Delta$
62.	A 26 yrs PG pregnant at 8 wks. Her hands were swollen with
05-	loss of the normally seen tendons & bony prominences.
	Which is not a possible diagnosis?
	a- Malnutrition
#	b- Renal insufficiency 6- PET
	d- Heart failure
	e- Liver cell failure
64-	Which is expected with that fetal position (star) at the onset
	of second stage of labor
	a- Normal labor
	a- Normal labor 6- Prolonged labor 7-7-7-6
	e-Precipitate labor
	d- Ulterine inertia
	e- None of the above

65- Which of the following don't contribute in the causation of such condition (star)

- a- Anthropoid pelvis
- b- Android pelvis
- e- DOP
- -d- Maternal kyphosis
- e- MIGR

66. Which is the incorrect statement regarding that technique

- for delivery of aftercoming head of breach (star)
- a- The aim is to deliver the fetus while maintaining full flexion of fetal head
- b- The technique is not initiated until moulding of the head
- c- Traction is not applied on the fetal body
- d- That technique could be aided by suprapubic pr
- e- The technique is abandoned in the current obst practice

7- Which is the incorrect statement regarding that fetal malpresentation

- a- CS is the most suitable method for delivery
- b- There is a risk of associated CFMF (hydroly)
 c- There is a risk of associated contracted pelvis
- de less common with PTL & Salinor.
- Is more common with twins

68- A case of obstructed labor with ant shoulder wedged behind the SP. Which is not a risk factor for such a case?

- a- Previous history of such event

- d- Macrosomia
- e- Anencephaly

69. That blotted graph during labor is useful in all the following except

- a. Early detection of abnormal progress of labor
- b- Estimation of rate of cervical dilatation
- e- Calculation of Bishop score
- d- Determining the need for augmentation, of labor
- e- Early diagnosis of obstructed labor

70- Kefaya keda, why?

- v- Ta3ban
- v= (ia3an
- z- Na3san

1- This cervix (star) is discovered at 20 wks preg, best ttt is

- a- Cryo-coagulation
- b- Laser coagulation
- c- Diathermy
- d- Cone biopsy
- e None of the above

2- That condition in the cervix, the true statement is

- a- Normally found in pregnancy
- b- Leads to lumbar backache (Sacal)
- e- Should be biopsied __oaluantimete
- d- Best treated by cone biopsy
- e- May lead to infertility

できいでしていた!

The ph is compared of chronic var discharge ?

Lockin-f.

3- Contraindication of that imaging study don't include

- a- Suspected blockage of Fallopian tubes
- b- Suspected acute PIDi Placer Juguist & organis
- c- Suspected intrauterine pregnancy
- d- Premenstrual phase -o Thick endown Polse -Ve (es. 1)
- e- During active bleeding from the aterus ____ extra 1/2:10 /1.

4- DES may lead to all the following except

- a- Patulous internal os
- b- T-shaped uterus
- c- Ectopic pregnancy
- d- Vaginal adenosis 一世 5分 元
- E- Septate merus

5- All the following regarding this test are true except

- a- Should be repeated in high risk population yearly
- b- Suspicious results should be confirmed by colposcopy
- Smear is only taken from the ectocervix (2 Port for 1)
- d- Is to be done for postcoital bleeding
- e- Suspicious specimen is related to HPV 16, 18

6- The result of this test have shown dysplastic cells occupying the deep 1/3 of the epithelium. What is your diagnosis

- Ja- CIN I
- b- CIM 2
- e- CIN 3
- d- CIN 4

7- This stain is done for all of these except

- a- Sexually active female with HPV 6, 11
- b- Sexually active female with HPV 16, 18
- c- Sexually active female esp smokers
- 'd- After subtotal hysterectomy
- _e Routinely in pregnancy

Cides

Sulleion.

8- Which is not associating such organism

- a- Vulval itching
- b- Redness of the vulva
- c- Odorless discharge
- d= Strawberry vaginal walls
- e- Dyspareunia

9- Which is the incorrect statement for that organism

- a- If may be discovered in the vaginal fluid of asymptomatic women
- be It flourish with increased vaginal acidity
- c- Usually associated with malodorous discharge
- d- May be associated with dysuria
- e- Vaginal discharge is typically frothy

10- The incorrect statement regarding that test

- a- Normally the epith of ectoex stains brown
- b- Vag wall stains brown during child bearing period
- c- Areas of columnar metaplasia don't stain brown -- c-
- d= Areas that don't stain brown are definitely malig -
- e- It is complementary test during colposcopy

11- All about this figure is true except

- त्रभी It is seen at mideyele
- b- Indicates estrogenic effect
- c- Positive in PCO at day 21 of cycle
- d- Turns --ve at day 18 of cycle
- e Indicates viscid mucoid cervix

12- This condition leads to all of these except

- a- Recurrent breech presentation
- b- Habitual abortion
- _d Ectopic pregnancy
- e- Menorrhagia -- to

13- All are true about such condition, except a Pelviabdominal swelling is mainly uterine b- Presents as Iry amenorrhea c- The commonest presentation is by retention of urine d- May lead to infertility if ttt is delayed —D Adhesions e- Needs surgical intervention	19-Regarding this swelling a It is the commonest cyst in the vulva b- It is present in the anterior part of labium majus (Post > c- Is mainly caused by obstruction of the duet by infection d- Should be excised if the patient is above 30 years e- The main symptom is throbbing pain
a- May be associated with uterus bicornus bicollus b- May be associated with horse show kidney c- Spasmodic dysmenorrhea could occur — A-Presents by Iry amenorrhea e- Presents by dyspareunia	20- Regarding this swelling, all are true except a- The causative organism is mainly E-coli b- Leads to throbbing pain c- Should be drained d- Is associated with pruritus vulvae (Pain) e- May occur due to gonorrhea
26 wks. Which is not a possible association? a- Recurrent breech b- PTL c- Absent left kidney d- Menorrhagia es Normal pregnancy & normal labor	21- All the following about this lesion is true except a Caused by gonorrhea b- Caused by HPV 6,11 c- May occur in the cervix d- May be treated by chemical cautery e- May be associated with CIN
degree perineal tear. The arrow indicates a structure called a- Bulbo-spongiosus b- Ischio-cavernosus c- Perineal body d- Transverse perineal ms e- None of the abo.e	a- It is an ulcer of the ectocervix b- During pregnancy is best treated by conization c- Cauterization should be the first line therapy d- Pap smear is advisable before therapy e- Commonly it causes pain, dysparuenia and backache 23- Causes of that condition doesn't include
7- All the following statements about such trauma are true except a- May be associated with diabetic mother b- High forceps may lead to it c- Surgical repair is done layer by layer d- Intercourse is avoided for 2-3 years (>>>> +6.)	a- Normal pregnancy b- COC c- ERT d- Acute cervicitis — Silo Borr Psp Carcinoma of cervix

24- Such polyps, all are true except

b- May be due to fibroid

a- Are commonly due to chronic cervicitis

c- Should be removed & sent for histopathology

e- Leads to contact bleeding (Mucous Polyte)

e- May be corrected under regional anesthesia
= epidual

18- Such a condition may be associated with, except

- a- Obstructed labor
- b- Lest occipito-anterior
- JUGI
- d- Precipitate labor (Sages
- e- Ureteric injury

25- This pathology is seen by hysteroscopy, the following should be done, except:

- a- Adhesiolysis via hysteroscopy
- b- Insertion of a balloon
- c- Giving steroids
- --- d= Giving progesterone
 - e- Giving cyclic estrogen & progesterone

26- This discharge is associated with, except:

- a- Acidic ph
- b- COC
- c- Severe soreness in the vulva
- _d-' Good response to metronidazole ___ t> -
- e- Bleeding spots after scrapping

27- This discharge is associated with, except

- a- Typically occurs postmenstrual
- b- May be associated with malodor
- c- Vagina is strawberry
- -d= Cultured on Sabaroud's medium
- e- Its organism may be seen in normal females

28- The arrow shows a condition associated with

- a- Normal vagina
- b- Pruritus & dysparuenia in most cases
- c- Responds well to miconazole
- et- Alkaline vaginal ph
- e- Odorless discharge ? _ m/

29- This condition is associated with all, except

- a- Recurrent lower abdominal pain & discharge
- b- Infertility
- c- Ectopic pregnancy
- d- Chlamydia
- e-1117 6,11 1 (X) ; L.

30- All the following about this slide is true, except

- a. Tubes are better removed in infertile patients
- b- IUCD must be removed before treatment
- c- Suspected more with chlamydial infection
- I Leads to habitual abortion
- e- Leads to ectopic pregnancy



MO(0:0/Piny Chronic

31-1 Pdf to development of prolapse include all except

- a- Congenital weakness of supporting ligaments
- b- Postmenopausal atrophy
- c- Injury during childbirth
- d- Cervical polyps
- e= Ovarian tumors

31-2 Which is the correct statement for that condition

- a- Usually associated with acute local pain _____
- b- More common in nulliparas
- c- Not related to instrumental delivery Tix 'vo Cointing
- d- Not related to postmenopausal changes
- May include sacral backache

3? - Regarding this test all are true except

- a- It may reach up to 15 cm
- b- It turns -ve 3 days after ovulation in Mullerian agenesis
- Tve test means water cervical mucous
- d- +ve test depends on estrogen
- e= +ve test means ovulation

33- All the following is true regarding IVF & ET except : Wild Alect

- 2 Down regulation is done by gonadotrophins Colonsing holon of Controls
- b- Ovarian stimulation may be done by Gn RH
- c- More than one embryo is transferred at a time -- Paradous L

JINLUGE

- d- Success rate is inversely proportional to age
- e-May be done in bilateral cornual block bidica 11.5

34- Which is the incorrect statement regarding that syndrome

- a- It is associated with gonadal failure
- b- There is no secondary sexual ecc
- c- The total number of enromosomes is 45
- d- Estrogen replacement therapy may be used _______.
- E- The ovaries may be polycystic

31:- All these statements about this syndrome is correct except

- a- Usually occurs in obese patients
- b- Is associated with insulin resistance
- c- Follicles are mainly seen subcapsular
- FSII/LII ratio is 3:1
- e- May lead to habitual abortion

49- Engagement of the fetal head is documented when

- a. The fetal head is inside the pelvis
- b- The presenting part is just above the level of ischial spines
- c- The vertex is left OA
- -d- The biparietal diameter pass through the pelvic inlet
- e- The fetal skull is fully flexed

50- All of theses are signs of placental separation except

- a- Uterus comes more globular & harder
- b- Suprapubic bulge
- e- Shortening of the umbilical cord
- d- Gush of blood from the vagina
- e- The fundus raises upwards

51- First trimester abortion may be due to: (100 Co minor est)

- a- Inadequate estrogen production in -t. Porton, 5
- H- Chromosome abnormality of the fetus
- e- Incompetence of the internal cervical os
- d- Maternal diabetes
- e- Cytotoxic drugs

52- Cervical incompetence

- a- Is a common cause of 1st trimester abortion
- b. May be congenital
- e- Is associated with previous instrumental delivery /
- d- is diagnosed at follicular phase HSG
- e- is associated with painful abortion &

53- In ectopic pregnancy:

- a- Bleeding precedes pain (Sal'
 - .5- Shoulders tip pain is an important symptom
- c- The isthmus of the tube is the commonest site of implantation (Arr Pullo)
- Let The incidence is greater in women with IUCD
- e- Ultrasonic scan is of no help in diagnosis

54- Which is the incorrect statement for that case

- 12-HCG is doubled every 48 hours
- Laparoscopy is the traditional standard method to confirm diagnosis
- Les Tubal pregnancy commonly presents with abdominal pain
- , d-Medical conservative III is an option on its by Ulbs < 200
- History of previous salpingitis is relevant

y last technologymonianthis with intent own. No. b.

55- The following statements regarding V.mole are correct except

- a- Usually have female karyotype
- b- May be complicated by thyrotoxicosis
- c- May be complicated by ovarian cyst
- d=α-seto-protein is a good marker for disease post-evacuation size G
- e- Is more prevalent in the advanced age group

Indications for a methotrexate chemotherapy following evacuation of a VM usually include except

- a- a rise in HCG titers
- b- plateau HCG titers for 3 successive weeks
- c- failure of HCG titers to return to normal 8 wks after evacuation
- d- appearance of liver or brain metastases
- rapid disappearance of theca lutein cysts (which

57-35 yrs old, G3P2 presented at 8 wks with mild vag bleeding for 3 days. She passes fleshy tissues per vagina presented on the slide. Further investigations don't include 11. 000 0

- a- Chest X-ray
- 13- Amniocentesis
- é- Serum HCG levels
- d- Pelvic U/S
- 58- Correlate
 - a) P.previa with severe bleeding & immature fetus (4)
 - b) P.previa centralis accidentally discovered at 37 wks (3)
 - (1) Mild placental abruption, no fetal distress at 33 wks
- e) Placental edge 3 cm from ex internal os, presenting in labor
 - . Conservative treatment
 - 2. Albowance or trial of vaginal delivery
 - 3. Elective cesarean section
 - 4. Emergency cesarean section
 - 5. Cesarean hysterectomy

59- As regards P. previa, the following are true

- a) It is manifested by painless bleeding
- b) The initial lige, is usually fatal
- c) It may predispose to PPHge
- d) Its incidence is unaffected by parity
- e) Its incidence decreases with maternal age

60-	The following are risk factors for placental abruption except
	a) Smoking
	b) Folic acid deficiency
	c) Preeclampsia
	History of threatened abortion " " " " " " " .
	e) Previous placental abruption $\frac{1000}{1000}$
<i>~</i> 9 9	This hematoma was found on inspection of the maternal side
01- 1	of the placenta after delivery. The risk factors for this
	condition don't include
	$(1-1)[\cdot]$
	b- Smoking to
	fron deficiency anemia $4 \times 100 \times 5 \times 1000 \times 1000$
	d- Chronic hypertension
	e- External trauma to the abdomen
62-	A diagnosis of severe PET in 37 wks with BPr 160/110 is
,	supported by
	a- Urine output of 1000 ml/24 hrs
	b- That physical sign on the slide - D Pital 13 555 Ands
11	e Epigastric pain Contra So.
	d- BPr of 160/110 at 8 wks in the same gestation 🗡
	e- Λ parity $> 5 \gamma \Delta$
62.	A 26 yrs PG pregnant at 8 wks. Her hands were swollen with
05-	loss of the normally seen tendons & bony prominences.
	Which is not a possible diagnosis?
	a- Malnutrition
#	b- Renal insufficiency 6- PET
	d- Heart failure
	e- Liver cell failure
64-	Which is expected with that fetal position (star) at the onset
	of second stage of labor
	a- Normal labor
	a- Normal labor 6- Prolonged labor 7-7-7-6
	e-Precipitate labor
	d- Ulterine inertia
	e- None of the above

65- Which of the following don't contribute in the causation of such condition (star)

- a- Anthropoid pelvis
- b- Android pelvis
- e- DOP
- -d- Maternal kyphosis
- e- MIGR

66. Which is the incorrect statement regarding that technique

- for delivery of aftercoming head of breach (star)
- a- The aim is to deliver the fetus while maintaining full flexion of fetal head
- b- The technique is not initiated until moulding of the head
- c- Traction is not applied on the fetal body
- d- That technique could be aided by suprapubic pr
- e- The technique is abandoned in the current obst practice

7- Which is the incorrect statement regarding that fetal malpresentation

- a- CS is the most suitable method for delivery
- b- There is a risk of associated CFMF (hydroly)
 c- There is a risk of associated contracted pelvis
- de less common with PTL & Salinor.
- Is more common with twins

68- A case of obstructed labor with ant shoulder wedged behind the SP. Which is not a risk factor for such a case?

- a- Previous history of such event

- d- Macrosomia
- e- Anencephaly

69. That blotted graph during labor is useful in all the following except

- a. Early detection of abnormal progress of labor
- b- Estimation of rate of cervical dilatation
- e- Calculation of Bishop score
- d- Determining the need for augmentation, of labor
- e- Early diagnosis of obstructed labor

70- Kefaya keda, why?

- v- Ta3ban
- v= (ia3an
- z- Na3san